

Home > Industries > Farms, fishing and forestry > Agriculture > Livestock > Movement of livestock > Transporting livestock between cattle tick zones > How to identify cattle tick

How to identify cattle tick

Cattle tick

Comparing cattle ticks to other ticks

Use the following table to distinguish cattle ticks from bush ticks and paralysis ticks.



Bush tick

Legs:

- Dark red-brown
- First pair close to snout

Legs:

- Pale cream
- · Wide space between first pair and snout

Paralysis tick/Scrub tick



Legs:

- · First and last pair brown, others pale
- In V-shape line from snout down sides of body

Bush tick	Cattle tick	Paralysis tick/Scrub tick
Body:	Body:	Body:
• Oval	Oval to rectangular	 Pear-shaped to oval
Face:	Face:	Face:
 Broadly oval 	• Oval, wider at front	Oval, wider at rear
Snout:	Snout:	Snout:
 Short, wider at face 	 Short, straight 	Very long

Cattle ticks in different stages

The following image shows what cattle ticks look like during different phases of development.





The following close-up shows 2 engorged female cattle ticks.

Also consider...

- View a map of cattle tick free and infested zones (http://arcg.is/1U1pD2f).
- Contact <u>Biosecurity Queensland (https://www.daf.qld.gov.au/about-us/contact-us)</u> for more information on cattle ticks.
- Read more about <u>cattle ticks (https://www.business.qld.gov.au/industries/farms-fishing-forestry/agriculture/livestock/animal-welfare/pests-diseases-disorders/cattle-tick)</u>.

Last reviewed: 03 Mar 2017

Last updated: 03 Mar 2017

© The State of Queensland 1995–2018

Queensland Government (http://www.qld.gov.au/)